DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

Lake: NATHAN	POND	Lake Area (ha):	16.19
Town:	DIXVILLE	Maximum depth (m):	6.4
	Coos	Mean depth (m):	2.4
River Basin:		Volume (m³):	389500
	44°55'13" N	Relative depth:	1.4
Longitude:		Shore configuration:	1.00
	2018	Areal water load (m/yr):	46.80
	(m): 1300	Flushing rate (yr^{-1}) :	19.50
Watershed are		P retention coeff.:	0.37
% watershed p	oonded: 0.0	Lake type: natural	l w/dam

BIOLOGICAL:	23 February 1993	4 August 1992
DOM. PHYTOPLANKTON (% TOTAL) #1	MELOSIRA 95%	MELOSIRA 65%
#2		DINOBRYON 20%
#3		
PHYTOPLANKTON ABUNDANCE (cells/mL)		360
CHLOROPHYLL-A (μg/L)		2.54
DOM. ZOOPLANKTON (% TOTAL) #1	FILINIA 55%	DAPHNIA 27%
#2	KERATELLA 25%	CALANOID COPEPODS 23%
#3		POLYARTHRA 18%
ROTIFERS/LITER	91	. 35
MICROCRUSTACEA/LITER	14	61
ZOOPLANKTON ABUNDANCE (#/L)	105	96
VASCULAR PLANT ABUNDANCE		Scattered
SECCHI DISK TRANSPARENCY (m)		3.2
BOTTOM DISSOLVED OXYGEN (mg/L)	1.3	0.0
BACTERIA (E. coli, #/100 ml) #1		200
#2		
#3		

SUMMER THERMAL STRATIFICATION:

stratified

Depth of thermocline (m): 4.5 Hypolimnion volume (m³): None Anoxic volume (m³): 28000

CHEMICAL:	Lake: NATHAN POND Town: DIXVILLE				
	23 February 1993		4 August 1992		
DEPTH (m)	2.0	4.0	2.0		4.0
pH (units)	6.2	6.1	7.0		6.9
A.N.C. (Alkalinity)	14.1	15.5	14.2		14.3
NITRATE NITROGEN	0.27	0.26	< 0.02		< 0.02
TOTAL KJELDAHL NITROGEN	0.14	0.20	0.28		0.34
TOTAL PHOSPHORUS	0.011	0.012	0.017		0.014
CONDUCTIVITY (µmhos/cm)	43.9	46.0	38.4	,	39.9
APPARENT COLOR (cpu)	34	41	33		42
MAGNESIUM			0.75		
CALCIUM			5.0		
SODIUM			0.9		
POTASSIUM			0.49		
CHLORIDE	< 3	< 3	< 3		< 3
SULFATE	4	4	4		4
TN : TP	37	38	16		24
CALCITE SATURATION INDEX			2.4		


All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 1992

D.O.	S.D.	PLANT	CHL	TOTAL	CLASS
**	2	1	0	3	Oligo.

COMMENTS:

- 1. This is a remote pond, located up a walk-in trail north of Dixville Notch. It was surveyed jointly with the Fish and Game Department.
- 2. There were no camps on the pond, but there was a camp in the woods approximately 1/4 mile from the pond.
- 3. Aphanocapsa (30%) and Cryptomonas (15%) were the dominant genera of wholewater phytoplankton. Blue-greens (40%), Cryptomonads (25%) and greens (25%) were the dominant classes.



FIELD DATA SHEET

LAKE: NATHAN POND

DATE: 08/04/92

TOWN: DIXVILLE

WEATHER: OVERCAST, COOL


	(°C)	OXYGEN	SATURATION
0.1	17.4	8.5	88 %
1.0	16.6	7.9	81 %
2.0	15.5	7.7	78 %
3.0	14.7	5.4	53 %
4.0	12.2	0.0	0 %
5.0	8.0	0.0	0 %
6.0	6.5	0.0	0 %
	-		
	2.0 3.0 4.0 5.0	2.0 15.5 3.0 14.7 4.0 12.2 5.0 8.0	2.0 15.5 7.7 3.0 14.7 5.4 4.0 12.2 0.0 5.0 8.0 0.0

SECCHI DISK (m): 3.2 COMMENTS:

BOTTOM DEPTH (m): 6.2

TIME: 1115

*Dissolved oxygen values are in mg/L



AQUATIC PLANT SURVEY

LAKE: NATHAN POND TOWN: DIXVILLE DATE: 08/04/92

PLANT	NAME	ABUNDANCE	
GENERIC	COMMON		
Sagittaria	Arrowhead	Scattered	
Juncus	Rush	Sparse	
Potamogeton	Pondweed	Sparse	
Nymphoides cordatum	Floating heart	Sparse	
Nuphar	Yellow water lily	Sparse	
Nymphaea	White water lily	Sparse	
Sparganium	Bur reed	Scattered	
	100		
	GENERIC Sagittaria Juncus Potamogeton Nymphoides cordatum Nuphar Nymphaea	Sagittaria Arrowhead Juncus Rush Potamogeton Pondweed Nymphoides cordatum Floating heart Nuphar Yellow water lily Nymphaea White water lily	

OVERALL ABUNDANCE: Scattered

GENERAL OBSERVATIONS:

1. A beaver hut and wood duck box were observed.